

A positive answer to the Riemann hypothesis: A new result predicting the location of zeros

Zeraoulia Elhadj

Department of Mathematics, University of Tébessa, (12002), Algeria

E-mail: zeraoulia@mail.univ-tebessa.dz and zelhadj12@yahoo.fr

December 27, 2012

Abstract

In this paper, a positive answer to the Riemann hypothesis is given by using a new result predicting the exact location of zeros of the alternating zeta function on the critical strip.

Keywords: Riemann hypothesis, alternating zeta function, location of zeros.

AMS 2010 *Mathematics Subject Classification*. *Primary* 11M26, *Secondary* 11A41, 11N05.

1 Introduction

Let $s = \alpha + i\beta$ be a complex number. The complex Riemann zeta function is defined in the half-plane $\alpha > 1$ by the absolutely convergent series

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} \quad (1)$$

and in the whole complex plane $\mathbb{C}$ by analytic continuation. As shown by Riemann, the zeta function (1) extends to $\mathbb{C}$ as a meromorphic function with only a simple pole at $s = 1$ with residue 1. In [1] Riemann obtained an analytic formula for the number of primes up to a preassigned limit in terms

of the zeros of the zeta function (1). This principal result implies that natural primes are distributed as regularly as possible if the Riemann hypothesis is true.

Riemann Hypothesis. The nontrivial zeros of $\zeta(s)$ have real part equal to $\alpha = \frac{1}{2}$.

The Riemann hypothesis is probably the most important open problem in pure mathematics today [2]. The unsolved Riemann hypothesis is part of the Hilbert's eighth problem, along with the Goldbach conjecture. It is also one of the Clay mathematics institute millennium prize problems. This hypothesis has been checked to be true for the first 1500000000 solutions. However, a mathematical proof is not formulated since its formulation in 1859.

It is well known that the Riemann hypothesis is equivalent to the statement that all the zeros of the Dirichlet eta function (the alternating zeta function)

$$\eta(s) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^s} \quad (2)$$

falling in the critical strip $D = \{s = \alpha + i\beta \in \mathbb{C} \text{ with } 0 < \alpha < 1\}$ lie on the critical line $\alpha = \frac{1}{2}$. See [3-4-8, pp. 49]. We have $\eta(s) = (1 - 2^{1-s}) \zeta(s)$ and the series $\eta(s)$ given by (2) converges only for $s = \alpha + i\beta \in \mathbb{C}$ with $\alpha > 0$. The function $\eta(s)$ is a non-constant analytic for all $s = \alpha + i\beta \in \mathbb{C}$ with $\alpha > 0$.

2 Evaluating the possible values of α

The functional equation for $\eta(s)$ restricted to the critical strip D is given by

$$\begin{cases} \eta(s) = \varphi(s) \eta(1-s) \\ \varphi(s) = 2^{\frac{(1-2^{-1+s})}{1-2^s}} \pi^{s-1} \sin\left(\frac{\pi s}{2}\right) \Gamma(1-s) \end{cases} \quad (3)$$

Let $\eta(s) = x(s) + iy(s)$ and $\eta(1-s) = u(s) + iv(s)$ be the algebraic forms of $\eta(s)$ and $\eta(1-s)$ where

$$\left\{ \begin{array}{l} x(s) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^{\alpha}} \cos(\beta \ln n), y(s) = - \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^{\alpha}} \sin(\beta \ln n) \\ u(s) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^{1-\alpha}} \cos(\beta \ln n), v(s) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^{1-\alpha}} \sin(\beta \ln n) \end{array} \right. \quad (4)$$

Also, the function $\varphi(s)$ can be written as $\varphi(s) = \varphi_1(s) + i\varphi_2(s)$ and therefore it is given uniquely by the polar form $\varphi(s) = \rho(s) \exp(i\theta(s))$, where, $\rho(s) = \sqrt{\varphi_1^2(s) + \varphi_2^2(s)} \neq 0$ is the magnitude and $\theta(s) = \arg \varphi(s) \in \mathbb{R}$, is the argument of $\varphi(s)$. Here, we have $\varphi_1(s) = \rho(s) \cos \theta(s)$ and $\varphi_2(s) = \rho(s) \sin \theta(s)$ since such complex numbers are entirely determined by their modulus and angle. The fact that $\varphi(s)$ is analytic implies that $\theta(s)$ and $\rho(s)$ cannot be constant everywhere in any open subset of $\mathbb{C}$. However, this is possible in *some* open sets of D as we can see below.

Note that all the identities concerning arguments in this paper holds only modulo factors of 2π if the argument is being restricted to $(-\pi, \pi]$. To compute these values we can use function $\arctan 2$ defined as follow

$$\arctan 2(a, b) = \left\{ \begin{array}{ll} \arctan\left(\frac{b}{a}\right), & \text{if } a > 0 \\ \arctan\left(\frac{b}{a}\right) \pm \pi, & \text{if } a < 0 \\ \pm \frac{\pi}{2}, & \text{if } a = 0 \\ \text{undefined} & \text{if } b = 0, a = 0 \end{array} \right.$$

with principal values in the range $(-\pi, \pi]$. Here, $(a, b) = a + ib \in \mathbb{C}$ and the sign "−" corresponding to the case $b < 0$. The main reason here is that the statement concerning the argument of a product used between Equations (12) and (13) below assumes the argument as an *equivalence class* of numbers. If we need the argument as a single number one needs to fix a half-open interval of length 2π to which the argument belongs by definition, i.e, we choose $(-\pi, \pi]$ for the purpose of calculations in this paper. The angle associated with the product of two complex numbers $z_1 z_2$ is $\theta_1 + \theta_2$. Hence, all the equations involving arguments from and below equation (12) are calculated using addition modulo 2π . We simply write $\arg(z_1 z_2) = (\theta_1 + \theta_2) \pmod{2\pi}$ and take only the *principal value* of any angle.

The main result for the proof of the Riemann hypothesis is given as follow:

Theorem 1 *The complex number $s = \alpha + i\beta \in D$ is a solution of $\eta(s) = 0$ and $\alpha = \frac{1}{2}$ if and only if $\theta(s) \neq 0 \pmod{2\pi}$, $x(s) = u(s)$ and $y(s) = -v(s)$.*

Proof. We note that the assumptions of Theorem 1 are valid only for single $s \in D$ since all zeros of a non-constant analytic functions are isolated. Hence, in all what follow we deals only with a single value $s \in D$.

(1) Proving that if $s = \alpha + i\beta \in D$ satisfying $\theta(s) \neq 0 \pmod{2\pi}$, $x(s) = u(s)$ and $y(s) = -v(s)$, then s is a solution of $\eta(s) = 0$ with $\alpha = \frac{1}{2}$:

For this purpose, substituting the algebraic forms of $\eta(s)$ and $\eta(1-s)$ into equation (3) to obtain

$$\left\{ \begin{array}{l} x(s) = u(s) \varphi_1(s) - v(s) \varphi_2(s) \\ y(s) = u(s) \varphi_2(s) + v(s) \varphi_1(s) \\ \text{or} \\ \begin{pmatrix} x(s) \\ y(s) \end{pmatrix} = A(s) \begin{pmatrix} u(s) \\ v(s) \end{pmatrix} \\ A(s) = \rho(s) \begin{pmatrix} \cos \theta(s) & -\sin \theta(s) \\ \sin \theta(s) & \cos \theta(s) \end{pmatrix} \end{array} \right. \quad (5)$$

and the inverse transformation is given by

$$\left\{ \begin{array}{l} u(s) = \frac{x(s)}{\rho^2(s)} \varphi_1(s) + \frac{y(s)}{\rho^2(s)} \varphi_2(s) \\ v(s) = -\frac{x(s)}{\rho^2(s)} \varphi_2(s) + \frac{y(s)}{\rho^2(s)} \varphi_1(s) \end{array} \right. \quad (6)$$

By setting $z(s) = \frac{x(s)}{\rho(s)}$ and $w(s) = \frac{y(s)}{\rho(s)}$, we obtain a *rotation transformation* of the form

$$\left\{ \begin{array}{l} \begin{pmatrix} z(s) \\ w(s) \end{pmatrix} = B(s) \begin{pmatrix} u(s) \\ v(s) \end{pmatrix} \\ B(s) = \begin{pmatrix} \cos \theta(s) & -\sin \theta(s) \\ \sin \theta(s) & \cos \theta(s) \end{pmatrix} \end{array} \right. \quad (7)$$

The matrix $B(s)$ in (7) is invertible for all $s \in D$ since its determinant is 1. It is well known that a *non trivial rotation* must have a unique fixed point, its *rotocenter*. The rotation in (5) is non trivial if $\varphi(s) \neq 1$ (here we assumed that $\theta(s) \neq 0 \pmod{2\pi}$ in the second part of Theorem 1). The reason is that the trivial rotation corresponding to the identity matrix, in which no rotation takes place. The *fixed point* of the rotation in (7) must satisfies $(I_2 - B(s))(u(s), v(s)) = 0$ where I_2 is the 2×2 unit matrix. The determinant of the matrix $(I_2 - B(s))$ is $-2(\cos \theta(s) - 1)$ and it is not zero since $\theta(s) \neq 0 \pmod{2\pi}$, this means that s is a solution of $\eta(s) = 0$ by using (3).

For $s \in D$, the assumptions $x(s) = u(s)$ and $y(s) = -v(s)$ can be reformulated by using (5) and (6) as follow:

$$\begin{cases} x(s) = u(s) \varphi_1(s) - v(s) \varphi_2(s) = u(s) = x(s) \varphi_1(s) + y(s) \varphi_2(s) \\ y(s) = u(s) \varphi_2(s) + v(s) \varphi_1(s) = -v(s) = -(y(s) \varphi_1(s) - x(s) \varphi_2(s)) \end{cases}$$

that is,

$$\begin{cases} u(s) \varphi_1(s) - v(s) \varphi_2(s) = x(s) \varphi_1(s) + y(s) \varphi_2(s) \\ u(s) \varphi_2(s) + v(s) \varphi_1(s) = -(-x(s) \varphi_2(s) + y(s) \varphi_1(s)) \end{cases}$$

by replacing the values of $\varphi_1(s) = \rho(s) \cos \theta(s)$, $\varphi_2(s) = \rho(s) \sin \theta(s)$, and the values of $x(s), y(s), u(s)$ and $v(s)$ from (4), we obtain the following equation

$$\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^\alpha} (1 - n^{2\alpha-1} \rho^2(s)) \exp(i\beta \ln n) = 0 \quad (8)$$

At this stage assuming $\theta(s) \neq 0 \pmod{2\pi}$, $x(s) = u(s)$ and $y(s) = -v(s)$, then s is a root of $\eta(s) = 0$. Since, we speak about a single and isolated root s , the only solution of (8) is obtained when

$$\alpha = \frac{1}{2} \quad (9)$$

for some $\beta \in \mathbb{R}$. Indeed, for $\alpha = \frac{1}{2}$, we have $\rho(s) = 1$ by using direct calculations. See [9] for more details. In this case $s = \frac{1}{2} + i\beta$ is a root of equation (9).

(2) Proving that if $s = \alpha + i\beta \in D$ is a solution of $\eta(s) = 0$ with $\alpha = \frac{1}{2}$ then $\theta(s) \neq 0 \pmod{2\pi}$, $x(s) = u(s)$ and $y(s) = -v(s)$:

For this purpose, assume that $s \in D$ is a solution of $\eta(s) = 0$ with $\alpha = \frac{1}{2}$, then, trivially, we have $x(s) = u(s) = 0$ and $y(s) = -v(s) = 0$. Thus we only need to prove that $\theta(s) \neq 0 \pmod{2\pi}$. From the second equation of (3) we have

$$\varphi\left(\frac{1}{2} + i\beta\right) = \sqrt{2}\pi^{i\beta - \frac{1}{2}} \left(\frac{2^{i\beta - \frac{1}{2}} - 1}{2^{i\beta + \frac{1}{2}} - 1}\right) \left(\cosh \frac{1}{2}\pi\beta + i \sinh \frac{1}{2}\pi\beta\right) \Gamma\left(\frac{1}{2} - i\beta\right) \quad (10)$$

and by direct calculations, we get

$$\left\{ \begin{array}{l} \pi^{i\beta - \frac{1}{2}} = \frac{1}{\sqrt{\pi}} (\cos(\beta \ln \pi) + i \sin(\beta \ln \pi)) \\ \frac{2^{i\beta - \frac{1}{2}} - 1}{2^{i\beta + \frac{1}{2}} - 1} = \frac{3}{4} \frac{\cos(\beta \ln 2) - \frac{2}{3}\sqrt{2}}{\cos(\beta \ln 2) - \frac{3}{4}\sqrt{2}} - \frac{1}{4}i \frac{\sin(\beta \ln 2)}{\cos(\beta \ln 2) - \frac{3}{4}\sqrt{2}} \\ \Gamma\left(\frac{1}{2} - i\beta\right) = \frac{\sqrt{\pi} \exp(-i(2\vartheta(\beta) + \beta \ln 2\pi + \arctan(\tanh \frac{\pi\beta}{2})))}{\sqrt{\cosh \pi\beta}} \end{array} \right. \quad (11)$$

where $\vartheta(t)$ is the Riemann Siegel function [10] given by

$$\vartheta(t) = \arg\left(\Gamma\left(\frac{2it + 1}{4}\right)\right) - \frac{\log \pi}{2}t, t \in \mathbb{R}.$$

The argument is chosen such that a continuous function is obtained and $\vartheta(0) = 0$ holds, i.e., in the same way that the *principal branch* of the log Gamma function is defined. Also, the argument of $\Gamma(s)$ is well defined and harmonic on the plane with a cut along the negative real axis.

On one hand, the last formula of equation (12) can be proved easily for $z = \mu + it$ by using the following equations

$$\left\{ \begin{array}{l} \cos\left(\frac{\pi z}{2}\right) = \frac{1}{\sqrt{2}} \sqrt{\cosh(\pi t)} \exp(-i \arctan(\tanh(\frac{\pi t}{2}))) \\ \Gamma\left(\frac{1}{2} + i\frac{t}{2}\right) = \left|\Gamma\left(\frac{1}{4} + i\frac{t}{2}\right)\right| \exp(i(\vartheta(t) + \frac{t}{2} \ln \pi)) \\ \Gamma(z) \Gamma(z + \frac{1}{2}) = 2^{1-2z} \sqrt{\pi} \Gamma(2z) \\ \Gamma(z) \Gamma(1-z) = \frac{\pi}{\sin \pi z} \end{array} \right.$$

and then we get the formula by setting $z = \frac{1}{4} + i\frac{t}{2}$ and replacing t by $-\beta$. Indeed, we have

$$\Gamma(2z) = 2^{2z-1} \pi^{-\frac{1}{2}} \Gamma(z) \frac{\pi}{\cos(\pi z)} \frac{1}{\Gamma\left(\frac{1}{2} - z\right)}$$

For $z = \frac{1}{4} + i\frac{t}{2}, t \in \mathbb{R}$, we have

$$\Gamma\left(\frac{1}{2} + it\right) = \frac{2^{-\frac{1}{2}+it}\pi^{\frac{1}{2}}}{\cos\left(\pi\left(\frac{1}{4} + i\frac{t}{2}\right)\right)} \frac{\Gamma\left(\frac{1}{4} + i\frac{t}{2}\right)}{\Gamma\left(\frac{1}{4} - i\frac{t}{2}\right)}$$

Hence, we have

$$\Gamma\left(\frac{1}{2} + it\right) = \frac{2^{it}\pi^{\frac{1}{2}}\pi^{it}\exp(2i\vartheta(t))}{\sqrt{\cosh \pi t} \exp(-i \arctan(\tanh(\frac{\pi t}{2})))}$$

Thus, we obtain

$$\Gamma\left(\frac{1}{2} + it\right) = \frac{\sqrt{\pi} \exp(i(2\vartheta(t) + t \ln(2\pi) + \arctan(\tanh(\frac{\pi t}{2}))))}{\sqrt{\cosh(\pi t)}}$$

For $t = -\beta$, we have

$$\Gamma\left(\frac{1}{2} - i\beta\right) = \frac{\sqrt{\pi} \exp(-i(2\vartheta(\beta) + \beta \ln(2\pi) + \arctan(\tanh(\frac{\pi\beta}{2}))))}{\sqrt{\cosh(\pi\beta)}}$$

because the function $t \rightarrow \arctan(\tanh(\frac{\pi t}{2}))$ is odd and since the function $\vartheta(t)$ is also odd as shown in [10]. We note that other proof of the last equation of (12) uses the functional equation of the zeta function.

On the other hand, we can calculate the arguments of the quantities in (12) as follow

$$\left\{ \begin{array}{l} \arg(\sqrt{2}) = 0 \pmod{2\pi} \\ \arg\left(\pi^{i\beta-\frac{1}{2}}\right) = (\beta \ln \pi) \pmod{2\pi} \\ \arg\left(\frac{2^{i\beta-\frac{1}{2}}-1}{2^{i\beta+\frac{1}{2}}-1}\right) = \varpi(\beta) \pmod{2\pi} \\ \arg\left(\cosh \frac{\pi\beta}{2} + i \sinh \frac{\pi\beta}{2}\right) = \phi(\beta) \pmod{2\pi} \\ \arg\left(\Gamma\left(\frac{1}{2} - i\beta\right)\right) = \psi(\beta) \pmod{2\pi} \\ \varpi(\beta) = -\arctan\left(\frac{1}{3} \frac{\sin(\beta \ln 2)}{\cos(\beta \ln 2) - \frac{2}{3}\sqrt{2}}\right) \\ \phi(\beta) = \arctan\left(\tanh \frac{\pi\beta}{2}\right) \\ \psi(\beta) = -(2\vartheta(\beta) + \beta \ln 2\pi + \arctan(\tanh \frac{\pi\beta}{2})) \end{array} \right. \quad (12)$$

The arguments: $t \ln \pi$, $\varpi(t)$, $\phi(t)$ and $\psi(t)$ in (12) are well defined since the functions $\pi^{it-\frac{1}{2}}$, $\cosh(\frac{1}{2}\pi t) + i \sinh(\frac{1}{2}\pi t)$, $\frac{2^{i\beta-\frac{1}{2}}-1}{2^{i\beta+\frac{1}{2}}-1}$ and $\Gamma(\frac{1}{2}-it)$ are not zero for all $t \in \mathbb{R}$. In particular, this is true if $t = \beta = \text{Im}(s)$.

Since the complex argument of a product of two numbers is equal to the sum of their arguments, then from (12) and the remark following Equation (4) we have

$$\theta\left(\frac{1}{2} + i\beta\right) = (0 + \beta \ln \pi + \varpi + \phi + \psi) \pmod{2\pi} \quad (13)$$

Hence, we have

$$\begin{cases} \theta\left(\frac{1}{2} + i\beta\right) = 2(g(\beta) - \vartheta(\beta)) \pmod{2\pi} \\ g(\beta) = -\frac{1}{2}\left(\beta \ln 2 + \arctan\left(\frac{1}{3} \frac{\sin(\beta \ln 2)}{\cos(\beta \ln 2) - \frac{2}{3}\sqrt{2}}\right)\right) \end{cases} \quad (14)$$

Thus, from the first equation of (14) we get

$$\vartheta(\beta) = \left[g(\beta) - \frac{1}{2}\theta\left(\frac{1}{2} + i\beta\right)\right] \pmod{2\pi} \quad (15)$$

The only possible value of equations $\theta\left(\frac{1}{2} + i\beta\right) = 2k\pi$ is when $k = 0$. Indeed, assume that $\theta\left(\frac{1}{2} + i\beta\right) = 2k\pi$. We know from [10] that $\vartheta(\beta)$ is an odd function and $g(\beta)$ is also an odd function by direct calculations. Hence, from (15) we have $\vartheta(-\beta) = -g(\beta) - k\pi = -g(\beta) + k\pi$, that is, $k = 0$. As mentioned above, we must prove that $\theta\left(\frac{1}{2} + i\beta\right) \neq 0 \pmod{2\pi}$ if $\frac{1}{2} + i\beta$ is a root of $\eta(s) = 0$. We know that the function $\vartheta(\beta)$ has only three roots 0 and $\pm 17.8455995405\dots$. Hence, the equation $g(\beta) - \frac{1}{2}\theta\left(\frac{1}{2} + i\beta\right) = 0$ has only the same three roots, that is the equation $\theta\left(\frac{1}{2} + i\beta\right) = 2g(\beta)$ has only the same three roots. The only possible case to get $\theta\left(\frac{1}{2} + i\beta\right) = 0 \pmod{2\pi}$ is when $\beta = 0$ since we have $g(0) = 0$, $g(17.8455995405) = -4.8774\dots$ and $g(-17.8455995405) = 4.8774\dots$. However, the real $s = \frac{1}{2} + i0$ is not a root of $\eta(s) = 0$ (we have $\eta\left(\frac{1}{2}\right) \simeq 0.60440$). Hence, we have proved that if $s \in D$ is a solution of $\eta(s) = 0$ with $\alpha = \frac{1}{2}$, then $\theta(s) \neq 0 \pmod{2\pi}$. ■

Theorem 1 proves that the Riemann hypothesis is true for the function $\eta(s)$ if and only if $\theta(s) \neq 0 \pmod{2\pi}$, $x(s) = u(s)$ and $y(s) = -v(s)$ for every root $s \in D$ of the equation $\eta(s) = 0$. Hence, to prove that the Riemann hypothesis is true for the function $\eta(s)$, we need only a proof of

the second part in Theorem 1 since we have an equivalence between the two cases. Clearly, if $s \in D$ is a root of $\eta(s) = 0$, then $x(s) = u(s) = 0$ and $y(s) = -v(s) = 0$. Hence, we need only to prove that if s is a root of $\eta(s) = 0$, then $\theta(s) \neq 0 \pmod{2\pi}$.

For this purpose, let us define the following relation in $\mathbb{C}^2$.

$$\forall (s, q) \in \mathbb{C}^2 : sRq \Leftrightarrow \theta(s) \neq 0 \pmod{2\pi} \text{ and } \theta(q) \neq 0 \pmod{2\pi}$$

Clearly, this is not an equivalence relation if there exist $s \in \mathbb{C}$ such that $\theta(s) = 0 \pmod{2\pi}$ (for this point, reflexivity fails). Define the set $\Omega = \{s \in D : \theta(s) \neq 0 \pmod{2\pi}\}$ such that the relation is an equivalence relation. Indeed, we can verify the following properties:

- (a) Reflexivity: $\forall s \in \Omega, sRs \Leftrightarrow \theta(s) \neq 0 \pmod{2\pi}$.
- (b) Symmetry: $\forall (s, q) \in \Omega^2 : sRq \Leftrightarrow \theta(s) \neq 0 \pmod{2\pi} \text{ and } \theta(q) \neq 0 \pmod{2\pi} \Rightarrow qRs$.
- (c) Transitivity: $\forall (s, q, h) \in \Omega^3 : sRq \text{ and } qRh \Leftrightarrow \theta(s) \neq 0 \pmod{2\pi} \text{ and } \theta(q) \neq 0 \pmod{2\pi} \text{ and } \theta(h) \neq 0 \pmod{2\pi} \Rightarrow sRh$.

There is only one equivalence class for this relation and it is defined by

$$\bar{s} = \{q \in \Omega : \theta(s) \neq 0 \pmod{2\pi} \text{ and } \theta(q) \neq 0 \pmod{2\pi}\}$$

From Theorem 1, each $s \in \Omega$ is a root of $\eta(s) = 0$.

From the relation $\eta(s) = (1 - 2^{1-s})\zeta(s)$, we know that if $\beta > 0$, then the first non trivial zero of $\eta(s) = 0$ in D is *computable* by hand and its value is $s_1 = \frac{1}{2} + 14.134725142i\dots$. For this root, we have $\varphi(s_1) = 0.99262 + 0.12128i$ and $\theta(s_1) = (0.12158) \pmod{2\pi}$. Hence, the *representative* of this equivalence class is s_1 . This logic implies that if $s \in D$ is any root of $\eta(s) = 0$, then $\theta(s) \neq 0 \pmod{2\pi}$ since we have $\theta(s_1) \neq 0 \pmod{2\pi}$.

To complete the proof, we need to prove that all roots $s \in D$ of $\eta(s) = 0$ are located in Ω . Indeed, the converse of Theorem 1 can be stated as follow:

$(\eta(s) \neq 0) \text{ or } (\alpha \neq \frac{1}{2})$ if and only if $(\theta(s) \neq 0 \pmod{2\pi}) \text{ or } (x(s) \neq u(s)) \text{ or } (y(s) \neq -v(s))$.

The partial case (the complex number $s = \alpha + i\beta \in D$ is a solution of $\eta(\alpha + i\beta) = 0$ and $\alpha \neq \frac{1}{2}$) is include in the part $(\alpha \neq \frac{1}{2})$. In this case, $\theta(s) = 0 \pmod{2\pi} \Rightarrow (\eta(\alpha + i\beta) = 0 \text{ with } \alpha \neq \frac{1}{2}) \text{ or } (\eta(s) \neq 0)$. From the proof of Theorem 1, we have $\theta(s) \neq 0 \pmod{2\pi} \Rightarrow \eta(s) = 0$, then the converse of this case is $\eta(s) \neq 0 \Rightarrow \theta(s) = 0 \pmod{2\pi}$ and it is also true. Hence we have $\eta(s) \neq 0 \Rightarrow \theta(s) = 0 \pmod{2\pi} \Rightarrow (\eta(\alpha + i\beta) = 0 \text{ with } \alpha \neq \frac{1}{2}) \text{ or } (\eta(s) \neq 0)$, thus we conclude that $\theta(s) = 0 \pmod{2\pi}$ implies only

$\eta(s) \neq 0$. Hence, we get $\theta(s) = 0 \pmod{2\pi} \Leftrightarrow \eta(s) \neq 0$, that is, $\theta(s) \neq 0 \pmod{2\pi} \Leftrightarrow \eta(s) = 0$.

Finally, we can conclude that

Theorem 2 *The Riemann hypothesis is true, i.e., all the nontrivial zeros of $\zeta(s)$ have real part equal to $\alpha = \frac{1}{2}$.*

Proof. The fact that the Dirichlet eta function (2) have the same zeros as the zeta function (1) in the critical strip D , implies that all nontrivial zeros of $\zeta(s)$ have real part equal to $\alpha = \frac{1}{2}$. ■

At this end, the hope that primes are distributed as regularly as possible is now become a truth by this unique solution of the Riemann hypothesis. Also, all propositions which are known to be equivalent to or true under the Riemann hypothesis are now correct. Examples includes, growth of arithmetic functions, Lindelöf hypothesis and growth of the zeta function, large prime gap conjecture...etc.

Acknowledgment

We would like to thank many colleagues for their valuable comments improving the quality of this paper: Michel L. Lapidus (University of California), Andrew Odlyzko (University of Minnesota), Sergey. K. Sekatskii (Ecole Polytechnique Fédérale de Lausanne), Marek Wolf (Cardinal Stefan Wyszyński University, Poland), Chris King (University of Auckland, Australia), Jinhua Fei, Peter Braun and Craig Feinstein.

References

- [1] G. F. B. Riemann, *Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse*, Monatsber. Königl. Preuss. Akad. Wiss. Berlin, 671-680, Nov. 1859.
- [2] M. Sautoy, *The music of the primes: Searching to solve the greatest mystery in mathematics*, HarperCollins, New York, 2004.
- [3] H. M. Srivastava and J. Choi, *Series associated with the Zeta and related functions*, Kluwer Academic Publishers, Dordrecht, Boston, and London, 2001,
- [4] J. Sondow, *Zeros of the alternating zeta function on the line $\operatorname{Re}(s) = 1$* , Amer. Math. Monthly, vol. 110, pp. 435-437, 2003.

- [5] J. Havil, *The zeros of zeta.*” §16.6 in *Gamma: Exploring Euler’s constant*. Princeton, NJ: Princeton University Press, pp. 193-196, 2003,
- [6] J. Derbyshire, *Prime obsession: Bernhard Riemann and the greatest unsolved problem in mathematics*. New York: Penguin, 2004. , pp. 384.
- [7] J. P. Serre, *A Course in arithmetic*, Chapter VI. Springer-Verlag, 1973.
- [8] S. Choi, B. Rooney and A. Weirathmueller, *The Riemann hypothesis: A resource for the aficionado and virtuoso alike*, Springer-Verlag, 2007.
- [9] E. C. Titchmarsh, D. R. Heath-Brown, *Theory of the Riemann Zeta function*, Oxford Science Publications, 1987.
- [10] H. M. Edwards, *Riemann’s Zeta Function*, New York: Dover Publications, 1974.